



# IFS alternative operations impact

Vanessa Bailey

Jet Propulsion Laboratory, California Institute of Technology

June 11, 2019

© 2018 California Institute of Technology. Government sponsorship acknowledged. The research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. The decision to implement the WFIRST mission will not be finalized until NASA's completion of the National Environmental Policy Act (NEPA) process. This document is being made available for information purposes only.

All calculations are for SNR = 10 assuming  
SPC Phase B coronagraph table & 20190130 sensitivities  
V=5 star, planet at 240 mas

**MUF=1** integration time ON TARGET. No overheads or ref star

Delta mag at 730nm	IFS int [hr]	Amici int [hr]	2% NB int [hr]	2.5% NB int [hr]	5% NB int [hr]
<b>18.25 = req</b>	3.6	2.3	1.9	1.4	0.7
19	11	6.2	5.1	3.8	1.7
20	55	29	23	17	7
21	310	160	120	90	36
<b>21.5 = RV</b>	800	400	300	210	84


# MUF=1 SNR=10 integration time ON TARGET. No overheads or ref star

Delta mag at 730nm	IFS int [hr]	Amici int [hr]	2% NB int [hr]	2.5% NB int [hr]	5% NB int [hr]
<b>18.25 = req</b>	3.6	2.3	1.9	1.4	0.7
19	11	6.2	5.1	3.8	1.7
20	55	29	23	17	7
21	310	160	120	90	36
<b>21.5 = RV</b>	800	400	300	210	84

- Amici integration time is less than IFS
- 7x2% NB filters
  - Cannot detect RV planet.
  - Could meet requirement (TODO: calc/sum for all filters. Maybe: check REQ performance level?)
- 2x2.5% + 5% NB filter
  - Can probably detect RV planet (TODO: get actual dmags in proposed filters and recalc)





# IFS concept of operations

- Pre-image planet in Band 1 imaging to determine location
  - Use to schedule spectroscopy, minimum 1 week wait
- Dig the dark hole in IFS mode on a bright *reference* star
  - WFC using IFS itself
- Long integration on *reference* star for RDI library
- Acquire *target* star & align with coronagraph
- Long integration on *target* star
  - Keep WFC solution from reference star
- Return to *reference* star for WFC touchup & reference images



# Narrowband filter concept of operations


- Pre-image planet in Band 1 imaging to determine location
  - Use to schedule spectroscopy, minimum 1 week wait
- Dig the dark hole in **imaging mode** on a bright *reference* star
  - WFC uses science filter itself if  $\leq 3.5\%$ , otherwise sub-bands are needed
- Long integration on *reference* star for RDI library
- Acquire *target* star & align with coronagraph
- Long integration on *target* star
  - Keep WFC solution from reference star
- Return to *reference* star for WFC touchup & reference images
- Repeat procedure sequentially for each NB filter (  $\geq 3$  filters )
  - Minor consequence: temporal variability of speckles vs. planet are difficult to distinguish



# Prism concept of operations: part 1


- Pre-image planet in Band 1 imaging to determine location
  - Use to schedule spectroscopy, minimum 1 week wait
- Dig the dark hole in **imaging mode** on a bright *reference* star
  - WFC uses 3-3.5% sub-bands. Expected to meet 30hr WFC time requirement.
  - WFC needs full image information => no slit or prism in the path
  - **More filter slots needed** to new accommodate WFC filters
- Align slit over area of interest on *reference* star
  - Match slit location on target star
  - **Tricky.**
- Long integration on *reference* star for RDI library
- Spectrum of un-occulted *reference* star
  - Use FSM to place star in slit? Or satellite spots/streaks?

...



# Prism concept of operations: part 2

- Acquire *target* star & align with coronagraph
  - Remove slit and prism for alignment
- Re-Align slit over area of interest
- Long integration on target star
  - Keep WFC solution from reference star
- Spectrum of un-occulted *target* star
  - Use FSM to place star in slit? Or satellite spots/streaks?
- Return to reference star for WFC touchup & reference images
  - Repeat slit acquisition procedure each time



# Slit alignment tolerance

- 2nm wavelength calibration planet-to-slit alignment tolerance
- 0.5px =
  - < 1.5nm wavecal error over all Band 3
  - < 5% slit flux loss at red end of Band 3
  - **10mas**  $\sim$  1/15 slit width
- **PAM req. precision meets alignment needs**
  - **Initial** placement RMS = 10  $\mu$ m = 21 mas. Does **NOT** meet wavecal need
  - **Fine adjustment** RMS  $\sim$  1 $\mu$ m  $\sim$  2 mas. **DOES** meet wavecal need
    - **15 min settling time** required before fine adjustments
  - Bench deformations will move PAMS & DICAM < 3mas (worst case).
- **Other image motion sources TBD – help?**
  - What are req's on image drift relative to PAMs and DIAM due to motion of mirrors & bench? Image motion must be smaller than PAM precision



# Slit fine alignment adds ~30min overhead

- **Wait 15 min** for PAM to settle
- Determining slit position (*needs more thought...*)
  - **How to illuminate?** Out of band (see below)? Remove filter entirely? Satellite spots/streaks ( $<1E-7$ )? Nudge star off-center under mask?
  - Can we just fit apparent edges? Or do we need to cross-correlate with a pattern?
  - **15?? min** required to find & iterate on slit location. **Automated?**



575nm 3.3%  
 $10^{-7} \sim 5min \text{ exp}$

Halpha 1%

730 nm 3.0%

**730 nm 15%**

825 nm 3.3%



All calculations are for SNR = 10 assuming  
SPC Phase B coronagraph table & 20190130 sensitivities  
V=5 star, planet at 240 mas

June 10, 2019 version of Bijan's exposure time calculator

**MUF=1** integration time ON TARGET. No overheads or ref star


Delta mag at 730nm	IFS int [hr]	Amici int [hr]	2% NB int [hr]	2.5% NB int [hr]	5% NB int [hr]
<b>18.25 = req</b>	3.6	2.3	1.9	1.4	0.7
19	11	6.2	5.1	3.8	1.7
20	55	29	23	17	7
21	310	160	120	90	36
<b>21.5 = RV</b>	800	400	300	210	84

Amici extra overhead compared to IFS assuming OS6-like (4x2hr blocks on target + 1x2hr block on ref)  
- Target star:  $t/2 * 0.5\text{hr}$       - Reference star:  $t/4 * 0.5\text{hr}$   
- 400hr on target => extra 150hr overheads for slit alignment Still faster than IFS.


# Modeling of PSF distortion near IWA and OWA required

At red end, more of PSF emerges from behind occultor

=> **appears shifted**



$\Delta \sim 0.2 \lambda/D$   
 $\sim 10 \text{ mas}$   
 $\sim 2 \text{ nm}$



# Preliminary conclusions

- Slit+prism
  - likely to meet L2 requirement *if* OK to have  $R < 50$  at red end
  - likely to match or exceed IFS sensitivity for equal amount of *clock* time
    - target + reference + overheads
    - Slit alignment is tricky and time-consuming
    - Can slight alignment be automated?
  - Space for new WFC filters needed
- Narrowband filters
  - Simplified operations compared to prism
  - 7 x 2% filters may be able to meet L2 requirement but *can't do better*
    - TODO: evaluate with MUFs?
  - 2 x 2.5% + 1 x 5% filter can probably detect CH<sub>4</sub> in an RV planet in <1000hr
    - does NOT meet L2 b/c it does not span 15% bandwidth or achieve  $R=50$